arXiv:0903.1924v2 [math.RT] 16 May 2009

MUTATION-CLASSES OF DIAGRAMS VIA INFINITE GRAPHS

THILO HENRICH

Abstract. We give a complete description of the cluster-mutation classes of diagrams of Dynkin types $\mathbb{A}, \mathbb{B}, \mathbb{D}$ and of affine Dynkin types $\mathbb{B}^{(1)}, \mathbb{C}^{(1)}, \mathbb{D}^{(1)}$ via certain families of diagrams.

1. Introduction

In 2002 Fomin and Zelevinsky introduced the concept of cluster algebras [8], which has been elaborated considerably in recent years. The theory has connections to many important areas in mathematics such as dual canonical bases for quantum groups or total positivity for algebraic groups, but also Poisson geometry and Teichmüller theory. To get a good overview, we recommend [13]. Cluster algebras are not constructed as algebras given by generators and relations, but rather by an iterative process called mutation. Therefore understanding mutation is essential for the whole theory.

At each step of the mutation process one obtains a special kind of weighted graph called a diagram. The set of diagrams obtained via mutation from a given diagram is called a mutation class, and it is an important problem to determine the mutation class of a diagram. By a result of Buan and Reiten [5] it is known that the cardinality of the mutation class of a diagram whose underlying undirected graph is of (affine) Dynkin type is finite. It is natural to ask whether these finite mutation classes have an explicit combinatorial description.

In this paper we give a precise description of the mutation classes of diagrams whose underlying graph is of Dynkin type $\mathbb{A}, \mathbb{B}, \mathbb{D}$ or affine Dynkin type $\mathbb{B}^{(1)}, \mathbb{C}^{(1)}$, $\mathbb{D}^{(1)}$. This description is given in terms of certain families of diagrams defined via infinite graphs. Note that the result for type $\mathbb{A}$ is already well known by work of Buan and Vatne [7] as well as Seven [15]. Moreover there is some overlap with independently achieved results by Vatne [16] for type $\mathbb{D}$. Our result is the following (cf. sections 2 and 3 for precise definitions):

Theorem 1.1. *Let Γ be a connected diagram. Then Γ is of mutation type*

(1) $\mathbb{A}$ *if and only if it is an A diagram.*
(2) $\mathbb{B}$ *if and only if it is a B diagram.*
(3) $\mathbb{D}$ *if and only if it is a $D_\star$ diagram, where* $\star \in \{(\bigcirc, n)_{n\geq 3}, \square, \boxslash, \perp\}$.
(4) $\mathbb{B}^{(1)}$ *if and only if it is a $B_{\star,B}, B_{\star\wedge\star'}$ diagram, where* $\star \in \{(\bigcirc, n)_{n\geq 3}, \square, \boxslash, \perp\}$ *and* $\star' \in \{B, \overleftrightarrow{B}\}$, *or a $B_{\square\wedge\boxslash}$ diagram.*
(5) $\mathbb{C}^{(1)}$ *if and only if it is a $C_{B,B}$ or $C_{B\wedge B}$ diagram.*
(6) $\mathbb{D}^{(1)}$ *if and only if it is a $D_{\star,\star'}, D_{\star\vee\star'}, D_{\star\wedge\star'}$ diagram, where* $\star$, $\star' \in \{(\bigcirc, n)_{n\geq 3}, \square, \boxslash, \perp, \overleftrightarrow{\boxslash}\}$, *or a $D_{\boxtimes}$ diagram.*

This work was partially supported by the DFG project Bu-1866/1-2.

Let us briefly comment on the work which has been done concerning the general question of describing mutation classes. Following the classification of cluster algebras of finite type by Fomin and Zelevinsky [9], efforts were made to find an effective way of determining whether a given diagram is mutation equivalent to one with underlying graph of Dynkin type or not. Seven [15] gave an answer in terms of so called 'minimal forbidden sub-diagrams', while Barot [1] reduced the problem to the question of cyclical orientability. The former result gives a satisfying answer in the case of non simply-laced diagrams, but if the diagram is simply-laced and contains more than nine vertices, the result by Seven produces a checklist of more than two hundred single diagrams plus five families of diagrams. Still, this list of diagrams is very interesting since it coincides with the Happel-Vossieck list [11], which characterizes all minimal representation infinite algebras. These connections were investigated in [6]. It would be interesting to see whether the families of graphs produced in our paper occur in a different context as well.

One example for applications of our results is the derived equivalence classification of cluster-tilted algebras (i.e. the endomorphism algebras of cluster-tilting objects in the cluster-tilting category), cf. [3, 4]. This was done for type $\mathbb{A}$ by Buan and Vatne [7] and for type $\mathbb{A}^{(1)}$ by Bastian [2]. In this context we would also like to mention the results for type $\mathbb{D}$ by Ge, Lv and Zhang [10], whose approach via triangulations of polygons was inspired by work of Schiffler [14].

The methods applied in this paper are purely combinatorial. In section 2 we recall the definitions of the (affine) Dynkin diagrams occurring in Theorem 1.1 and the notion of diagram mutation. Section 3 then lists, for each (affine) Dynkin type, a certain number of infinite undirected graphs inducing the families of diagrams mentioned in the Theorem. The final section 4 contains the proof of Theorem 1.1.

2. (Affine) Dynkin Diagrams and Mutation of Diagrams

2.1. **Some (affine) Dynkin Diagrams.** Recall the definitions of the (affine) Dynkin diagrams mentioned in Theorem 1.1, which are

- the Dynkin diagrams of type $\mathbb{A}, \mathbb{B}, \mathbb{D}$:



- the affine Dynkin diagrams of type $\mathbb{B}^{(1)}, \mathbb{C}^{(1)}, \mathbb{D}^{(1)}$ (we also include type $\mathbb{A}^{(1)}$, which will be used in the definitions and proofs):



$B_n^{(1)}$



$C_n^{(1)}$



D_n



2.2. **Some Notations.** For any graph Γ, we denote by Γ_0 and Γ_1 the set of vertices respectively edges of Γ. The cardinality of Γ_0 is denoted by $|\Gamma_0|$ and for $x \in \Gamma_0$, $\deg(x)$ is the cardinality of the set of neighbours of x in the underlying undirected graph of Γ. If Γ is directed we also distinguish between $\deg^-(x)$, the number of neighbours y of x in Γ such that there exists an element $\{y \to x\}$ in Γ_1, and $\deg^+(x) := \deg(x) - \deg^-(x)$.

Given any subset $S \subseteq \Gamma_0$, the *induced sub-graph on* S is the sub-graph of Γ obtained from Γ by deleting all vertices in $\Gamma_0 \setminus S$ and all edges adjacent to these. A sub-graph $\Gamma' \subseteq \Gamma$ is called *full*, if Γ' is the induced sub-graph on Γ'_0. A directed graph Γ is called *cyclically oriented* if for all full sub-graphs $\Gamma' \subseteq \Gamma$ with underlying graph a cycle (i.e. equal to $A^{(1)}_{|\Gamma'_0|}$), Γ' is oriented clockwise or anti-clockwise.

For $\star$ any (affine) Dynkin type and Γ any directed graph, Γ is called a $\star$ graph if the underlying undirected graph of Γ is of type $\star$.

2.3. **Diagram mutation.** Let Γ be a finite directed graph whose edges are weighted with positive integers. Γ is called a *diagram* if the product of weights along any cycle is an integer which is the square of an integer. For any vertex $k \in \Gamma_0$, the *mutation of* Γ *in* k, denoted by μ_k, is the following transformation of Γ:

- the orientations of all edges incident to k are reversed, their weights remain unchanged.
- for any vertices i and j which are connected in Γ via a two-edge oriented path going through k (refer to the figure below for the rest of notation), the direction of the edge (i,j) in $\mu_k(\Gamma)$ and its weight c' are uniquely determined by the rule
$$\pm\sqrt{c} \pm \sqrt{c'} = \sqrt{ab}$$
where the sign before $\sqrt{c}$ (respectively before $\sqrt{c'}$) is $+$ if i, j, k form an oriented cycle in Γ (resp., in $\mu_k(\Gamma)$), and is $-$ otherwise. Here either c or c' can be equal to 0.



- the rest of the edges and their weights in Γ remain unchanged.

Note that the resulting weighted graph is a diagram again. Two diagrams Γ and Γ' related by a sequence of mutations $\mu_{i_1} \circ \mu_{i_2} \circ \ldots \circ \mu_{i_n}$, where $\{i_j\}_{1 \leq j \leq n} \subseteq \Gamma_0$, are called *mutation equivalent.* Since μ_k is involutive, i.e. $\mu_k^2(\Gamma) = \Gamma$, this defines an equivalence relation on the set of all diagrams. The equivalence classes are called *mutation classes.* The task of understanding these classes is not trivial at all. For example, we shall see that diagram i) below is neither mutation equivalent to diagram ii) nor to diagram iii), while ii) and iii) belong to the same mutation class.



It is well known that for diagrams with underlying undirected graph being a tree, the mutation class is independent of the particular orientation (cf. [9] Prop. 9.2). Hence we say that for $\star$ an (affine) Dynkin type other than $\mathbb{A}^{(1)}$, a diagram Γ is of *mutation type* $\star$ if it is mutation equivalent to a diagram with underlying undirected graph of type $\star$. Finally, a diagram is called *simply-laced* if all its edge weights are equal to one.

Remark 2.1. Let us stress that for the rest of this paper, the term 'diagram' will refer exclusively to a directed and weighted graph in the sense just described. Hence the word will never refer to an (affine) Dynkin diagram nor will there be any possibility for confusion with the common meaning of a diagram, which is that of an undirected graph.

3. The diagrams

The following notion will be involved in the definition of all diagrams for any type.

Definition 3.1. By a (∇, x) *graph* we refer to the infinite undirected graph



with uniquely determined vertex x. It is obtained by starting with an infinite binary tree and joining the two children of each vertex by an edge. We also write ∇ graph for short.

Before we give the diagrams for the particular (affine) Dynkin types, note that the notation used in parts (4) and (6) of Theorem 1.1 is slightly imprecise, as it suggests the existence of some families of diagrams which will not be defined (e.g. $B_{\perp\wedge\perp}$ or $D_{\perp\wedge\perp}$) . Let it be understood that these families are empty.

3.1. The diagrams for type $\mathbb{A}$.

Definition 3.2. A is the infinite undirected graph obtained by the following construction. Let (∇_1, x_1) and (∇_2, x_2) be two disjoint ∇ graphs. Identify x_1 and x_2.



An A *diagram* is a cyclically oriented simply-laced diagram with underlying undirected graph equal to a connected and full sub-graph of A.

3.2. The diagrams for type $\mathbb{B}$.

Definition 3.3. Let B be the infinite undirected weighted graph obtained by the following construction. Let (∇, x) be a ∇ graph. Assign weight two to both edges having x as one endpoint and weight one to all others.



A B *diagram* is a cyclically oriented diagram with underlying undirected weighted graph equal to a connected and full sub-graph of B containing the vertex x.

3.3. The diagrams for type $\mathbb{D}$.

Definition 3.4. 1) For $n \in \mathbb{N}_{\geq 3}$ let $D_{(\bigcirc,n)}$ denote the infinite undirected graph which is obtained by the following construction. Let $\{a_1, .., a_n\}$ be the set of vertices of an $A_{n-1}^{(1)}$ graph labeled clockwise, $(\nabla_i, x_i)_{1\leq i\leq n}$ be ∇ graphs. For all $1 \leq i \leq n$ join x_i to both a_i and a_{i+1} by single edges, where a_{n+1} denotes the vertex a_1.



The sub-graph of $D_{(\bigcirc,n)}$ induced on $\{a_i\}_{1\leq i\leq n}$ be denoted by $(\bigcirc, n)$. A $D_{(\bigcirc,n)}$ *diagram* Γ is a cyclically oriented simply-laced diagram with underlying undirected graph equal to a connected and full sub-graph of $D_{(\bigcirc,n)}$ containing the vertices of $(\bigcirc, n)$ such that $|\Gamma_0| \geq 5$.

2) $D_{\Box}$ is the infinite undirected graph obtained by the following construction. Let a_1, a_2 be two single vertices and $(\nabla_i, x_i)_{1\leq i\leq 2}$ be two ∇ graphs. Join each of x_1, x_2 to each of a_1, a_2 by a single edge.

By $\Box$ we denote the sub-graph of $D_{\Box}$ induced on $\{x_1, a_1, x_2, a_2\}$. A $D_{\Box}$ *diagram* is a cyclically oriented simply-laced diagram with underlying undirected graph equal to a connected and full sub-graph of $D_{\Box}$ containing the vertices of $\Box$.







3) Let $D_{\boxslash}$ denote the infinite undirected graph which is obtained by inserting one edge between x_1 and x_2 in $D_{\Box}$.

The sub-graph of $D_{\boxslash}$ induced by $\{a_i, x_i\}_{1\leq i\leq 2}$ is denoted by $\boxslash$. A $D_{\boxslash}$ *diagram* is a cyclically oriented simply-laced diagram with underlying undirected graph equal to a connected and full sub-graph of $D_{\boxslash}$ containing the vertices of $\boxslash$.

4) $D_{\perp}$ is the infinite undirected graph obtained by deleting the ∇ graph attached to x_2 in $D_{\Box}$ including the vertex x_2.

By $\perp$ we denote the sub-graph of $D_{\perp}$ induced on $\{a_1, a_2, x_1\}$. A $D_{\perp}$ *diagram* is a cyclically oriented simply-laced diagram with underlying undirected graph equal to a connected and full sub-graph of $D_{\perp}$ containing the vertices of $\perp$.

Remark 3.5. Note that the families of $D_\star$ diagrams, where $\star \in \big\{(\bigcirc, n)_{n\geq 3}, \Box, \boxslash, \perp\big\}$, are mutually disjoint. Indeed, the crucial point is that:

- a $D_{(\bigcirc,4)}$ diagram can always be distinguished from a $D_{\Box}$ diagram. This is because the first one is required to obtain at least five vertices, so it always contains a sub-graph of the form *i)* as shown below. But a $D_{\Box}$ diagram never does.
- a $D_{(\bigcirc,3)}$ diagram Γ can always be distinguished from a $D_{\boxslash}$ diagram even if two of $\{x_i\}_{1\leq i\leq 3}$ are absent (cf. *ii)* for the general case). This is because the first one always contains a sub-graph of the form *iii)* since $|\Gamma_0| \geq 5$, while the second one never does.

i)



ii)



iii)



3.4. **The diagrams for type $\mathbb{B}^{(1)}$.**

Definition 3.6. For any $\star \in \{(\bigcirc, n)_{n\geq 3}, \square, \boxslash, \perp\}$ a $B_{\star,B}$ *diagram* is a diagram obtained by the following construction. Let Γ be a $D_\star$ diagram, Γ' be a B diagram and fix the notations of definitions 3.4 and 3.3, where we add primes to the labels used in 3.3. Choose a vertex y of Γ which is not labeled a_i and $y' \in \Gamma'_0 \setminus \{x'\}$ such that the following is true:

- if $y \in \star_0$ then all neighbours of y are contained in $\star_0$.
- y' is of degree at most two, as is y if $y \notin \star_0$.

Then identify y and y'.

Given a $B_{\star,B}$ diagram $\bar{\Gamma}$, let $\omega_{\bar{\Gamma}}$ be the shortest of all paths in the underlying undirected graph starting in any of $x_i \in \Gamma_0$ and ending in $x' \in \Gamma'_0$ (again cf. the notation of definitions 3.2 and 3.3). Then the width $w(\bar{\Gamma})$ of $\bar{\Gamma}$ is defined as the length of $\omega_{\bar{\Gamma}}$ minus one.

Example 3.7. We give two examples, the first showing a $B_{(\bigcirc,4),B}$ diagram of width two, the second a $B_{\boxslash,B}$ of width zero.

i)



ii)



Remark 3.8. Note that for any $\star \in \{(\bigcirc, n)_{n\geq 3}, \square, \boxslash, \perp\}$ there are usually many different pairs of $D_\star$ and B diagrams which yield the same $B_{\star,B}$ diagram (for certain choices of vertices y and y' as described in the construction of definition 3.6). Cf. *i)* in the example above, where 'splitting' at any vertex in the horizontal line of x_2 yields such a pair. Let us stress that we are not interested in the combinatorics of such 'splittings', but only in the disjoint classes of $B_{\star,B}$ diagrams, where $\star \in \{(\bigcirc, n)_{n\geq 3}, \square, \boxslash, \perp\}$, themselves (cf. remarks 3.13 and 3.17).

Definition 3.9. 1) Let $B_{\star\wedge B}$, where $\star \in \{\square, \boxslash\}$, be the infinite undirected weighted graph obtained by the following construction. Delete the ∇ graph attached to x_1 in $D_\star$ without removing x_1. Assign weight two to all edges with endpoint x_1 and weight one to all others.

By $\star \wedge B$ we denote the sub-graph of $B_{\star\wedge B}$ induced on $\{x_i, a_i\}_{1\leq i\leq 2}$. A $B_{\star\wedge B}$ *diagram* is a cyclically oriented diagram with underlying undirected weighted graph equal to a connected and full sub-graph of $B_{\star\wedge B}$ containing the vertices of $\star \wedge B$.







2) $B_{\boxslash\wedge\square}$ is the infinite undirected graph obtained by joining a_1 and a_2 in $B_{\square\wedge B}$ by two edges of weight one.

The sub-graph of $B_{\boxslash\wedge\square}$ induced on $\{x_i, a_i\}_{1\leq i\leq 2}$ is denoted $\boxslash \wedge \square$. A $B_{\boxslash\wedge\square}$ *diagram* is a cyclically oriented diagram with underlying undirected weighted graph equal to a connected and full sub-graph of $B_{\boxslash\wedge\square}$ containing the vertices of $\boxslash \wedge \square$.

3) Let $B_{(\bigcirc,n)\wedge B}$ be the infinite undirected weighted graph obtained by the following construction. Delete the ∇ graph attached to x_1 in $D_{(\bigcirc,n)}$ without removing the vertex itself. Assign weight two to both edges with endpoint x_1 and weight one to all others.

For $\star \in \{B, \overleftrightarrow{B}\}$ the sub-graph of $B_{(\bigcirc,n)\wedge B}$ induced on $\{a_i\}_{1\leq i\leq n} \cup \{x_1\}$ is denoted by $(\bigcirc, n)\wedge\star$. A $B_{(\bigcirc,n)\wedge\star}$ *diagram* is a diagram Γ with underlying undirected weighted graph equal to a connected and full sub-graph of $B_{(\bigcirc,n)\wedge B}$ containing the vertices of $(\bigcirc, n) \wedge \star$ such that the following conditions are satisfied:

- if $\star = B$, then Γ is cyclically oriented.
- if $\star = \overleftrightarrow{B}$, let e denote the edge connecting a_1 and a_2. Then the graph $(\Gamma_0, \Gamma_1 \setminus e)$ is cyclically oriented, while the sub-graph induced on $\{a_i\}_{1\leq i\leq n}$ is not.

Example 3.10. The first example is a $B_{(\bigcirc,3)\wedge B}$, the second a $B_{(\bigcirc,3)\wedge\overleftrightarrow{B}}$ diagram. We indicate possible enlargements by dots.



3.5. The diagrams for type $\mathbb{C}^{(1)}$.

Definition 3.11. A $C_{B,B}$ *diagram* is a diagram obtained by the following construction. Let Γ, Γ' be two B diagrams and fix the notations of definition 3.3 (where we add primes to the labels used in 3.3 for the vertices of Γ'). Choose $y \in \Gamma_0 \setminus \{x\}$ and $y' \in \Gamma'_0 \setminus \{x'\}$ such that both vertices are of degree at most two. Identify y and y'.

For a $C_{B,B}$ diagram $\bar{\Gamma}$, let $\omega_{\bar{\Gamma}}$ be the shortest of all paths in the underlying undirected graph from x to x'. Then the width $w(\bar{\Gamma})$ of $\bar{\Gamma}$ is defined as the length of $\omega_{\bar{\Gamma}}$ minus two.

Example 3.12. The first example shows a $C_{B,B}$ diagram of width zero, the second a $C_{B,B}$ diagram of width one.



Remark 3.13. There are usually many different pairs of two B diagrams which yield the same $C_{B,B}$ diagram. Cf. *ii)* in the example above, where 'splitting' at any vertex in the right-hand triangle except at the lowest one yields such a pair. We are not interested in the combinatorics of such 'splittings', but only in the class of $C_{B,B}$ diagrams itself (cf. remarks 3.8 and 3.17).

Definition 3.14. Let $C_{B\wedge B}$ be the infinite undirected weighted graph obtained by the following construction. Let $(\nabla, \bullet)$ be a ∇ graph, x_1, x_2 two single vertices. Join each of x_1, x_2 to $\bullet$ by an edge of weight two, join both of them by an edge of weight four and assign weight one to all edges.



The sub-graph of $C_{B\wedge B}$ induced on $\{x_1, x_2, \bullet\}$ is denoted $B \wedge B$. A $C_{B\wedge B}$ *diagram* is a cyclically oriented diagram with underlying undirected weighted graph equal to a connected and full sub-graph of B containing the vertices of $B \wedge B$.

3.6. **The diagrams for type $\mathbb{D}^{(1)}$.**

Definition 3.15. A $D_{\star,\star'}$ *diagram*, where $\star, \star' \in \{(\bigcirc, n)_{n\geq 3}, \Box, \boxslash, \perp\}$, is a diagram obtained by the following construction. Let Γ be $D_\star$ diagram, Γ' be a $D_{\star'}$ diagram and fix the notations of definition 3.4 (where we add primes to the labels used in 3.4 for the vertices of Γ'). Choose a vertex y of Γ which is not labeled a_i and a vertex y' of Γ' which is not labeled a'_j such that the following conditions are fulfilled:

- if $y \in \star$ then all neighbours of y belong to $\star_0$ and $y' \notin \star'_0$.
- if $y' \in \star'$ then all neighbours of y' belong to $\star'_0$ and $y \notin \star_0$.
- any of y, y' not contained in $\star_0$ respectively $\star'_0$ is of degree at most two.

Finally, identify y and y'.

Given a $D_{\star,\star'}$ diagram $\bar{\Gamma}$, let $\omega_{\bar{\Gamma}}$ be the shortest of all paths in the underlying undirected graph from any of $x_i \in \Gamma_0$ to any of $x'_j \in \Gamma'_0$. Then the width $w(\bar{\Gamma})$ of $\bar{\Gamma}$ is defined as follows:

- if $(\star, \star') = ((\bigcirc, n), \bigcirc, m))$ then $w(\bar{\Gamma})$ is equal to the length of $\omega_{\bar{\Gamma}}$.
- otherwise $w(\bar{\Gamma})$ is equal to the length of $\omega_{\bar{\Gamma}}$ minus one.

Example 3.16. We give three examples. The first is a $D_{(\bigcirc,4),\perp}$ diagram of width two, the second is a $D_{\boxslash,\boxslash}$ diagram of width zero, the last is a $D_{(\bigcirc,3),(\bigcirc,3)}$ diagram of width zero.

i)



ii)



iii)



Remark 3.17. Note that there are usually many different pairs of $D_\star$ and $D_{\star'}$ diagrams, where $\star, \star' \in \{(\bigcirc, n)_{n\geq 3}, \Box, \boxslash, \perp\}$, which yield the same $D_{\star,\star'}$ diagram (for certain choices of vertices y and y' as described in the construction of definition 3.15). Cf. *i)* in the example above, where 'splitting' at any vertex in the horizontal row of x_2 or in x'_1 yields such a pair. Let us stress that we are not interested in the combinatorics of such 'splittings', but only in the disjoint classes of $D_{\star,\star'}$ diagrams, where $\star, \star' \in \{(\bigcirc, n)_{n\geq 3}, \Box, \boxslash, \perp\}$, themselves (cf. also remarks 3.8 and 3.13).

Definition 3.18. $D_{\star\vee\star'}$ is the infinite graph obtained by the following construction:

- if $(\star, \star') \neq ((\bigcirc, n), (\bigcirc, m))$ then delete the ∇ graphs attached to x_1, x'_1 in $D_\star$ respectively $D_{\star'}$ without deleting x_1, x'_1 themselves. Identify x_1 and x'_1 and denote this vertex by $\bullet$.
- if $(\star, \star') = ((\bigcirc, n), (\bigcirc, m))$ then delete the ∇ graphs attached to x_1, x_2 in $D_{(\bigcirc,n)}$ respectively x'_1, x'_2 in $D_{(\bigcirc,m)}$ without deleting these four vertices themselves. Identify the first and second vertex of the following pairs: $(x_1, a'_1), (x_2, a'_3), (a_1, x'_1), (a_3, x'_2)$. Finally identify a_2 and a'_2 and denote this vertex by $\bullet$.

The sub-graph of $D_{\star\vee\star'}$ induced on the vertices of $\star \cup \star'$ after identification is denoted by $\star \vee \star'$. A $D_{\star\vee\star'}$ *diagram* is a cyclically oriented simply-laced diagram Γ with underlying undirected graph equal to a connected and full sub-graph of $D_{\star\vee\star'}$ containing the vertices of $\star \vee \star'$. In case of $(\star, \star') = ((\bigcirc, n), (\bigcirc, m))$, we additionally require that $|\Gamma_0| \geq 6$.

Example 3.19. We give three examples, the first being a $D_{(\bigcirc,4)\vee(\bigcirc,3)}$, the second a $D_{(\bigcirc,3)\vee\boxslash}$ and the last a $D_{\boxslash\vee\perp}$ diagram.



Definition 3.20. 1) Let $n \in \mathbb{N}_{\geq 3}$. Then $D_{(\bigcirc,n)\wedge\Box}$ is the infinite undirected graph obtained by the following construction. Let $\{a_1, .., a_{n-1}\}$ be an A_{n-1} graph labeled linearly, $(\nabla_i, x_i)_{1\leq i\leq n-2}$ be ∇ graphs, $\{\bullet_j, a'_j\}_{1\leq j\leq 2}$ be a $A_4^{(1)}$ graph such that a_1 and a_2 are not connected. For all $1 \leq i \leq n-2$ join x_i to both a_i and a_{i+1} by single edges and identify $\bullet_1$ with a_{n-1} and $\bullet_2$ with a_1.





By $(\bigcirc, n)\wedge\square$ we denote the sub-graph of $D_{(\bigcirc,n)\wedge\square}$ induced on $\{a_i\}_{2\leq i\leq n-2}\cup\{\bullet_j, a'_j\}_{1\leq j\leq 2}$. A $D_{(\bigcirc,n)\wedge\square}$ *diagram* Γ is a simply-laced diagram with underlying undirected graph equal to a connected and full sub-graph of $D_{(\bigcirc,n)\wedge\square}$ containing the vertices of $(\bigcirc, n)\wedge\square$ such that the following condition is satisfied: each full sub-graph Γ' of Γ with underlying undirected graph a cycle is cyclically oriented, except for the case where Γ'_0 equals $\{a_i\}_{2\leq i\leq n-2}\cup\{\bullet_1, \bullet_2, a'_2\}$.

2) For $n\in\mathbb{N}_{\geq 3}$ let $D_{(\bigcirc,n)\wedge\boxslash}$ denote the infinite undirected graph obtained by joining $\bullet_1, \bullet_2$ in $D_{(\bigcirc,n+1),\square}$ by a single edge.

For any of $\star\in\{\boxslash, \overset{\leftrightarrow}{\boxslash}\}$, the sub-graph of $D_{(\bigcirc,n)\wedge\boxslash}$ induced on $\{a_i\}_{2\leq i\leq n-1}\cup\{\bullet_j, a'_j\}_{1\leq j\leq 2}$ is denoted by $(\bigcirc, n)\wedge\star$. A $D_{(\bigcirc,n)\wedge\star}$ *diagram* is a simply-laced diagram Γ with underlying undirected graph equal to a connected and full sub-graph of $D_{(\bigcirc,n)\wedge\boxslash}$ containing the vertices of $(\bigcirc, n)\wedge\star$ such that the following conditions are satisfied.

- If $\star=\boxslash$, then Γ is cyclically oriented.
- If $\star=\overset{\leftrightarrow}{\boxslash}$, let e denote the edge between $\bullet_1$ and $\bullet_2$. Then the graph $(\Gamma_0, \Gamma_1\setminus e)$ is cyclically oriented, while the sub-graph induced on $\{\bullet_1, \bullet_2\}\cup\{a_i\}_{2\leq i\leq n-1}$ is not.

Example 3.21. The first example shows a $D_{(\bigcirc,3)\wedge\boxslash}$, the second is a $D_{(\bigcirc,3)\wedge\overset{\leftrightarrow}{\boxslash}}$ diagram. We indicate possible enlargements by dots.



Definition 3.22. 1) $D_{\square\wedge\square}$ is the infinite graph obtained by the following construction. Take the induced sub-graph on $\square\subset D_{\square}$ and let $(\nabla, \bullet)$ be a ∇ graph. Connect both x_1, x_2 to $\bullet$.

By $\square\wedge\square$ we denote sub-graph of $D_{\square\wedge\square}$ induced on the vertices $\{a_i, x_i\}_{1\leq i\leq 2}\cup\{\bullet\}$. A $D_{\square\wedge\square}$ *diagram* is a simply-laced diagram Γ with underlying undirected graph equal to a connected and full sub-graph of $D_{\square\wedge\square}$ containing the vertices of $\square\wedge\square$ such that the following is fulfilled.

- For $1\leq i\leq 2$, the sub-graph induced on the set of vertices $\{x_1, \bullet, x_2, a_i\}$ is cyclically oriented.
- The sub-graph induced induced on $\Gamma_0\setminus\{a_i, x_i\}_{1\leq i\leq 2}$ is cyclically oriented.

2) By $D_{\boxslash\wedge\boxslash}$ we denote the infinite graph obtained by the following inserting two edges between x_1 and x_2 in $D_{\square\wedge\square}$.

The sub-graph of $D_{\boxslash\wedge\boxslash}$ induced on the vertices of $\{a_i, x_i\}_{1\leq i\leq 2}\cup\{\bullet\}$ is denoted by $\boxslash\wedge\boxslash$. A $D_{\boxslash\wedge\boxslash}$ *diagram* is a cyclically oriented simply-laced diagram with underlying undirected graph equal to a connected and full sub-graph of $D_{\boxslash\wedge\boxslash}$ containing the vertices of $\boxslash\wedge\boxslash$.

Example 3.23. The first example depicts a $D_{\square\wedge\square}$, the second a $D_{\boxslash\wedge\boxslash}$. We also include a $D_{(\bigcirc,3)\wedge\square}$ diagram to make the reader aware of the subtle difference in the definitions involved. Again, possible enlargements are indicated by dots.



Definition 3.24. $D_{\boxtimes}$ is the infinite graph obtained by the following construction. Let $\{a_i\}_{1\leq i\leq 4}$ be an $A_4^{(1)}$ graph , (∇, x_1) a ∇ graph. Join each of $\{a_i\}_{1\leq i\leq 4}$ with x_1 via one edge.



The sub-graph of $D_{\boxtimes}$ induced on the vertices of $\{a_i\}_{1\leq i\leq 4} \cup \{x_1\}$ is denoted by $\boxtimes$. A $D_{\boxtimes}$ *diagram* is a cyclically oriented simply-laced diagram with underlying undirected graph equal to a connected and full sub-graph of $D_{\boxtimes}$ containing the vertices of $\boxtimes$.

Remark 3.25. Note that the families of $D_{\boxtimes}$,$D_{\star,\star'}$, $D_{\star\vee\star'}$ and $D_{\star\wedge\star'}$ diagrams with $\star,\star' \in \left\{(\bigcirc,m)_{m\geq 3},\square,\boxslash,\perp,\overleftrightarrow{\boxslash}\right\}$ are mutually disjoint. Indeed, the crucial point is that any $D_{(\bigcirc,3)\vee(\bigcirc,3)}$ diagram Γ (see *i)* below for the general case) may be distinguished from a $D_{\boxtimes}$ diagram. This is because the first mentioned always contains an induced underlying sub-graph of the form *ii)* since $|\Gamma_0| \geq 6$. But a $D_{\boxtimes}$ diagram never does.



4. Proof

The idea for the proof is the same for all types. In the first step we prove that the set of families of graphs defined for a particular type is invariant under mutation. The second step then applies these results to show that any diagram belonging to one of the families defined for a particular type indeed is an element of that type's mutation class.

Mutation is a local concept, hence the first step mentioned above involves a detailed examination of the effect of diagram mutation at a given vertex (depending on the local situation). The results are presented in Facts 4.1. − 4.7. We shall sketch the proofs for Facts 4.1 and 4.2. only, leaving the rest to the reader. To get a good feeling for the statements, we recommend the applet by Keller [12].

4.1. **Type $\mathbb{A}$.**

Fact 4.1. Let Γ be an A diagram, k any vertex of Γ. Then $\mu_k(\Gamma)$ is an A diagram.

Proof. All cases to be considered are encoded by

$$\begin{tikzcd}
2 \arrow[rr] & & 3 \arrow[dl] & & 2 \arrow[dr] & & 3 \arrow[dd] \\
& k \arrow[ul] \arrow[dr] & {} \arrow[rr, leftrightarrow, "\mu_k"] & & {} & k \arrow[dl] \arrow[ur] & \\
1 \arrow[ur] & & 4 \arrow[ll] & & 1 \arrow[uu] & & 4 \arrow[ul]
\end{tikzcd}$$

□

Theorem. 1.1 (1) *Let Γ be a connected diagram. Then Γ is of mutation type $\mathbb{A}$ if and only if it is an A diagram.*

Proof. Since any diagram with underlying graph of Dynkin type $\mathbb{A}$ is itself an A diagram, one direction follows from Fact 4.1. For the other direction, we proceed by induction on $|\Gamma_0|$.

For $|\Gamma_0| = 1$ there is nothing to show. So assume Γ has $n+1$ vertices. If no vertex belongs to a cycle we are done. Else choose a vertex y such that the induced sub-graph Γ' on $\Gamma_0 \setminus \{y\}$ is connected and apply the inductive assumption to Γ'. Denote by μ a series of mutations transforming Γ' into a diagram with underlying undirected graph A_n. Then $\mu(\Gamma)$ has underlying undirected graph A_n with y joined to some of its vertices, but it is also an A diagram by Fact 4.1. If y is connected to only one vertex, then the underlying graph of $\mu(\Gamma)$ must already be A_{n+1}. Assuming that y is connected to more than two vertices of A_n, we label the vertices of A_n linearly and choose the smallest three which are joined to y, say $i < j < k$

$$\begin{tikzcd}
& & y \arrow[dl, dash] \arrow[d, dash] \arrow[dr, dash] & & \\
1 \arrow[r, dash, dashed] & i \arrow[r, dash, dashed] & j \arrow[r, dash, dashed] & k \arrow[r, dash, dashed] & n
\end{tikzcd}$$

Since A contains no cycles of length $m \geq 3$, it follows that $j = i+1$ and $k = i+2$. But then $\mu(\Gamma)$ is obviously no A diagram. Therefore y is connected to at most two vertices of $\mu(\Gamma)$, which have to be neighbours as we just proved. Thus mutation in y yields a diagram with underlying undirected graph A_{n+1}. □

4.2. **Type $\mathbb{B}$.**

Fact 4.2. Let Γ be an B diagram, k any vertex of Γ. Then $\mu_k(\Gamma)$ is an B diagram.

Proof. All cases to be considered beyond that dealt with in Fact 4.1 are encoded by

$$\begin{tikzcd}
x \arrow[ddr, "2"] & & & x \arrow[dd, "2"] & \\
& {} \arrow[rr, leftrightarrow, "\mu_k"] & & {} & \\
\circ \arrow[uu, "2"] & \circ \arrow[l] & & \circ \arrow[r] & \circ \arrow[uul, "2"']
\end{tikzcd}
\qquad
\begin{tikzcd}
x \arrow[rr, "2"] & & 3 \arrow[dl] & & x \arrow[dr, "2"] & & 3 \arrow[dd] \\
& k \arrow[ul, "2"] \arrow[dr] & {} \arrow[rr, leftrightarrow, "\mu_k"] & & {} & k \arrow[dl] \arrow[ur] & \\
1 \arrow[ur] & & 2 \arrow[ll] & & 1 \arrow[uu, "2"] & & 2 \arrow[ul]
\end{tikzcd}$$

□

Theorem. 1.1 (2) *Let Γ be a connected diagram. Then Γ is of mutation type $\mathbb{B}$ if and only if it is a B diagram.*

Proof. Use the same arguments as in the proof for type $\mathbb{A}$ with the appropriate replacements. For the inductive step we additionally assume that $y \neq x$, which is always possible. $\square$

4.3. **Type $\mathbb{D}$.** The following result is an easy exercise in diagram mutation (cf. Definition 3.4 for notations).

Fact 4.3. Let Γ be a $D_\star$ diagram for $\star \in \big\{(\bigcirc, n)_{n\geq 3}, \square, \boxslash, \perp\big\}$ and k be any vertex of Γ.

(1) If $k \notin \star_0$, then $\mu_k(\Gamma)$ is a $D_\star$ diagram.

Else assume

(2) $\star = (\bigcirc, n)$. Then mutation in k yields a
  (a) $D_\square$ diagram, if $n = 3$, $k \in \{a_i\}_{1\leq i\leq 3}$ and $x_j \in \Gamma_0$, where x_j is the uniquely determined vertex among $\{x_i\}_{1\leq i\leq 3}$ not connected to k.
  (b) $D_\perp$ diagram if $n = 3$, $k \in \{a_i\}_{1\leq i\leq 3}$ and $x_j \notin \Gamma_0$, x_j as before.
  (c) $D_{(\bigcirc,n-1)}$ diagram if $n > 3$, $k \in \{a_i\}_{1\leq i\leq n}$.
  (d) $D_{(\bigcirc,n+1)}$ diagram if $k \in \{x_i\}_{1\leq i\leq n}$.

(3) $\star = \square$. Then mutating in k yields a
  (a) $D_{\bigcirc,3}$ diagram if $|\Gamma_0| > 4$, $k \in \{x_1, x_2\}$.
  (b) $D_\boxslash$ diagram if $k \in \{a_1, a_2\}$, or both $|\Gamma_0| = 4$ and $k \in \{x_1, x_2\}$

(4) $\star = \boxslash$. Then mutating in k yields a
  (a) $D_\square$ diagram if $k \in \{a_1, a_2\}$.
  (b) $D_\boxslash$ diagram if $k = x_i$ for some $i \in \{1, 2\}$ and k has a neighbour $y \in (\nabla_i)_0$ such that the induced sub-diagram of Γ on $\{y, x_1, x_2\}$ carries non-linear orientation and a $D_\perp$ diagram else.

(5) $\star = \perp$.
  (a) If $k \in \perp \setminus \{x\}$, mutation in x yields a $D_\perp$ diagram.
  (b) Else $x = k$ and $\mu_k(\Gamma)$ is a
    (i) $D_\boxslash$ diagram if there exists a neighbour y of x such both $\{y, x, a_i\}_{1\leq i\leq 2}$ carry linear orientation.
    (ii) $D_\perp$ diagram if x is of degree three and there exists a neighbour $y \notin \perp_0$ of x such both $\{y, x, a_i\}_{1\leq i\leq 2}$ carry non-linear orientation.
    (iii) $D_{(\bigcirc,3)}$ diagram else.

Before proving the Theorem for type $\mathbb{D}$, let us state the following Lemma.

Lemma 4.4. *Let $n \geq 3$ and C be a cyclically oriented simply-laced diagram with underlying undirected graph $A^{(1)}_{n-1}$, whose vertices are labeled counterclockwise by $\{1, ..., n\}$, Γ' be any diagram, $y \in \Gamma'_0$. Denote by Γ the diagram obtained by adding the arrows $\{y \to 1\}$ and $\{n \to y\}$ of weight one. If $\mu := \mu_{n-2} \circ \cdots \circ \mu_2 \circ \mu_1$, then $\mu(\Gamma) \setminus \Gamma'$ has underlying undirected graph D_n and $\{1 \to y\}$ is the only element of*

Γ_1 *between* $(\mu(\Gamma)\setminus\Gamma')_0$ *and* Γ'_0 *in any direction. Moreover the induced sub-graph on* $\{y\}\cup\{1,..,n\}$ *is simply-laced.*



Proof. The statement follows by induction using the following two facts





□

Now we can easily prove:

Theorem. 1.1 (3) *Let* Γ *be a connected diagram. Then* Γ *is of mutation type* $\mathbb{D}$ *if and only if it is a* $D_\star$ *diagram, where* $\star\in\big\{(\bigcirc,n)_{n\geq 3},\Box,\boxslash,\perp\big\}$.

Proof. Any diagram with underlying undirected graph of Dynkin type $\mathbb{D}$ is a $D_\perp$ diagram, hence Fact 4.3 implies that any diagram of mutation type $\mathbb{D}$ is a $D_\star$ diagram for some $\star\in\big\{(\bigcirc,n)_{n\geq 3},\Box,\boxslash,\perp\big\}$.

To show the other direction, first assume that $|\Gamma_0|=4$. Then $\star\notin\{(\bigcirc,n)_{n\geq 3}\}$ by definition. Moreover, by Fact 4.3 (3) (b) and (4) (b) we may assume that $\star=\perp$. Thus Γ has underlying undirected graph D_4.

For $|\Gamma_0|>4$ we first show that any $D_\star$ diagram with $\star\in\{\perp,\boxslash,\Box\}$ is mutation equivalent to a $D_{(\bigcirc,m)}$ diagram, $m\leq|\Gamma_0|$. By Fact 4.3 (5) (b) we may restrict to $\star\in\{\boxslash,\Box\}$; here we use that if we are in the situation of sub-case (5) (b) (ii), then $\mu_{a_1}(\Gamma)$ is as described in one of the other sub-cases of (5) (b). Now (4) (a) of the same Fact allows for further restriction to $\star=\Box$, hence we may apply (3) (a).

Thus we may assume that Γ is a $D_{(\bigcirc,m)}$ diagram, $m\leq|\Gamma_0|$. Set $n:=|\Gamma_0|$ and apply Fact 4.3 (2) (d) until Γ is a cycle on n vertices. We infer from Lemma 4.4 that Γ is mutation equivalent to a diagram with underlying undirected graph D_n. □

4.4. **Type $\mathbb{B}^{(1)}$.** Besides the difficulty of keeping track of notations (cf. sub-section *3.4*), the following Fact is a straightforward exercise in diagram mutation.

Fact 4.5. The following is true.

(1) Let Γ be a $B_{\star,B}$ diagram, where $\star\in\{(\bigcirc,n)_{n\geq 3},\Box,\boxslash,\perp\}$. Recall that the endpoints of ω_Γ are labeled x_i for some i and x'. Let Γ_1 denote the connected component of the sub-graph of Γ induced on $\Gamma_0\setminus\{x'\}$ which is uniquely determined by containing x_i. Then Γ_1 is a $D_\star$ diagram and a full connected sub-diagram of Γ. For k any vertex of Γ we distinguish between two cases. If $k\in(\Gamma_1)_0$ then $\mu_k(\Gamma_1)$ is a $D_{\star'}$ diagram for some $\star'\in\big\{(\bigcirc,n)_{n\geq n},\Box,\boxslash,\perp\big\}$ by Fact 4.3. Otherwise we simply set $\star'=\star$. Then the following is true:

   (a) if $w(\Gamma)>0$, then $\mu_k(\Gamma)$ is a $B_{\star',B}$ diagram and $|w(\mu_k(\Gamma))-w(\Gamma)|\leq 1$
   .

- (b) if $w(\Gamma) = 0$ and $k \neq x_i$, then $\mu_k(\Gamma)$ is a $B_{\star',B}$ diagram and $w(\Gamma) \leq w(\mu_k(\Gamma)) \leq w(\Gamma) + 1$.
- (c) otherwise $w(\Gamma) = 0$ and $k = x_i$ which is a neighbour of x'. Assume that $\star$ equals
    - (i) $(\bigcirc, n)$. Then $\mu_k(\Gamma)$ is a $B_{(\bigcirc,n+1)\wedge B}$ diagram.
    - (ii) $\square$. Then $\mu_k(\Gamma)$ is a $B_{(\bigcirc,3)\wedge B}$ diagram.
    - (iii) $\boxslash$. Then $\mu_k(\Gamma)$ is a
        - (A) $B_{\boxslash\wedge B}$ diagram if $\{x', x_1, x_2\}$ carries non-linear orientation.
        - (B) $B_{\boxslash,B}$ diagram of width zero if $\{x', x_1, x_2\}$ carries linear orientation and $\deg(x) = 5$.
        - (C) $B_{\perp,B}$ diagram of width zero else.
    - (iv) $\perp$. Then $\mu_k(\Gamma)$ is a
        - (A) $B_{\boxslash\wedge B}$ diagram if both $\{x', x, a_i\}_{1\leq i\leq 2}$ carry linear orientation.
        - (B) $B_{\boxslash,B}$ diagram of width zero if both $\{x, x', a_i\}_{1\leq i\leq 2}$ carry non-linear orientation and $\deg(x) = 4$.
        - (C) $B_{\perp,B}$ diagram of width zero if both $\{x, x', a_i\}_{1\leq i\leq 2}$ carry non-linear orientation and $\deg(x) = 3$.
        - (D) $B_{(\bigcirc,3)\wedge B}$ diagram else.

(2) Let Γ be a $B_{(\bigcirc,n)\wedge\star}$ diagram, $\star \in \{B, \overleftrightarrow{B}\}$.

- (a) If $k \notin (\bigcirc, n) \wedge B$ then $\mu_k(\Gamma)$ is a $B_{(\bigcirc,n)\wedge\star}$ diagram.
- (b) If $k = x_1$ then $\mu_k(\Gamma)$ is a $B_{(\bigcirc,n)\wedge\overleftrightarrow{B}}$ diagram if $\star = B$ and a $B_{(\bigcirc,n)\wedge B}$ diagram if $\star = \overleftrightarrow{B}$.
- (c) If $k \in \{x_i\}_{i\neq 1}$ then $\mu_k(\Gamma)$ is a $B_{(\bigcirc,n+1)\wedge\star}$ diagram.
- (d) If $k \in \{a_i\}_{i\neq 1,2}$ and $n > 3$, then $\mu_k(\Gamma)$ is a $B_{(\bigcirc,n-1)\wedge\star}$ diagram.
- (e) If $n = 3$ and $k = a_3$, then $\mu_k(\Gamma)$ is a $B_{\square\wedge B}$ diagram if $\star = B$. Else $\star = \overleftrightarrow{B}$ and μ_k yields a $B_{\boxslash\wedge\square}$ diagram.
- (f) Else $k \in \{a_1, a_2\}$. Assume $\star$ equals
    - (i) $\overleftrightarrow{B}$. Then $\mu_k(\Gamma)$ is a $B_{(\bigcirc,n)\wedge\overleftrightarrow{B}}$ diagram.
    - (ii) B. Then
        - (A) for $n > 3$, $\mu_k(\Gamma)$ is a $B_{(\bigcirc,n-1),B}$ diagram of width zero.
        - (B) for $n = 3$, $\mu_k(\Gamma)$ is a $B_{\square,B}$ diagram of width zero if $x_i \in \Gamma_0$, where x_i is the uniquely determined vertex among $\{x_j\}_{1\leq j\leq 3}$ not connected to k and a $B_{\perp,B}$ diagram of width zero else.

(3) Let Γ be a $B_{\star\wedge\star'}$ diagram, $\star, \star' \in \{\square, \boxslash, B\}$.

- (a) If $k \notin (\star \wedge \star')_0$ then $\mu_k(\Gamma)$ is a $B_{\star\wedge\star'}$ diagram.
- (b) Else $k \in (\star \wedge \star')_0$.
    - (i) If $k = x_2$ then $\mu_k(\Gamma)$ is a
        - (A) $B_{(\bigcirc,3)\wedge B}$ diagram if $\star \wedge \star' = \square \wedge B$.
        - (B) $B_{\boxslash,B}$ diagram of width zero if $\star \wedge \star' = \boxslash \wedge B$ and there exists a neighbour $y \notin (\boxslash \wedge B)_0$ of x_2 such that $\{x_1, x_2, y\}$ carries non-linear orientation.
        - (C) $B_{\perp,B}$ of width zero if $\star \wedge \star' = \boxslash \wedge B$ and $y \notin \Gamma_0$, y as before.
        - (D) $B_{(\bigcirc,3)\wedge\overleftrightarrow{B}}$ diagram if $\star \wedge \star' = \boxslash \wedge \square$.

- (ii) If $k = x_1$ then $\mu_k(\Gamma)$ is a
  - (A) $B_{(\boxslash\wedge\Box)}$ diagram if $\star \wedge \star' = \Box \wedge B$ and vice versa.
  - (B) $B_{(\boxslash\wedge B)}$ diagram else.
- (iii) Else $k \in \{a_1, a_2\}$ and $\mu_k(\Gamma)$
  - (A) $B_{(\boxslash\wedge B)}$ diagram if $\star \wedge \star' = \Box \wedge B$ and vice versa.
  - (B) $B_{(\boxslash\wedge\Box)}$ diagram else.

The idea for the difficult part of the proof of Theorem 1.1 (4) is to apply the above result to show that any diagram belonging to one of the families defined in subsection 3.4 is of mutation type $\mathbb{B}^{(1)}$. This is done by reducing successively the cases to be considered.

Theorem. 1.1 (4) *Let Γ be a connected diagram. Then Γ is of mutation type $\mathbb{B}^{(1)}$ if and only if it is a $B_{\star,B}, B_{\star\wedge\star'}$ diagram, where $\star \in \big\{(\bigcirc, n)_{n\geq 3}, \Box, \boxslash, \perp\big\}$ and $\star' \in \{B, \overleftrightarrow{B}\}$, or a $B_{\Box\wedge\boxslash}$ diagram.*

Proof. Any diagram with underlying undirected graph of Dynkin type $\mathbb{B}^{(1)}$ is a $B_{\perp,B}$ diagram, hence Fact 4.5 implies that any diagram of mutation type $\mathbb{B}^{(1)}$ is a $B_{\star,B}, B_{\star\wedge\star'}$ diagram, where $\star \in \big\{(\bigcirc, n)_{n\geq 3}, \Box, \boxslash, \perp\big\}$ and $\star' \in \{B, \overleftrightarrow{B}\}$, or a $B_{\Box\wedge\boxslash}$ diagram.

To show the other direction, first assume that $|\Gamma_0| = 4$. This restricts the possibilities for Γ to $B_{\star\wedge\star'}$ diagrams, where $\star \in \big\{(\bigcirc, 3), \Box, \boxslash\big\}$ and $\star' \in \{B, \overleftrightarrow{B}\}$, or $B_{\boxslash\wedge\Box}$, or $B_{\perp,B}$ diagrams. Note that the last mentioned is a $B_4^{(1)}$ graph already. Thus Fact 4.5 (3) (b) (i) (C) implies that any $B_{\boxslash\wedge\Box}$ diagram is of mutation type $\mathbb{B}^{(1)}$. Hence we need only deal with $B_{\star\wedge\star'}$ diagrams, where $\star \in \big\{(\bigcirc, 3), \Box, \boxslash\big\}$ and $\star' \in \{B, \overleftrightarrow{B}\}$. Now Fact 4.5 (2) (e) allows to reduce to the case of $B_{\Box\wedge B}$ and $B_{\boxslash\wedge\Box}$ diagrams. But any of these are mutation equivalent to a $B_{\boxslash\wedge B}$ diagram by (3) (b) (iii) (A) and (3) (b) (ii) (B) respectively. Since we already proved $B_{\boxslash\wedge B}$ diagrams to be of mutation type $\mathbb{B}^{(1)}$, this finishes the proof for $|\Gamma_0| = 4$.

For $|\Gamma_0| > 4$ we apply the results of Fact 4.5 in order to reduce to the case where Γ is a $B_{(\bigcirc,m)\wedge B}$ diagram for some $m \leq |\Gamma_0| - 2$. First note that any $B_{\star,B}$ diagram, where $\star \in \big\{(\bigcirc, m)_{m\geq 3}, \Box, \boxslash, \perp\big\}$, is mutation equivalent to a $B_{\star',B}$ diagram, where $\star' \in \big\{(\bigcirc, m)_{m\geq 3}, \Box, \boxslash, \perp\big\}$, of width zero. Next Fact 4.5 (1) (c) reduces to the case of $B_{\star\wedge\star'}$ diagrams, where $\star \in \big\{(\bigcirc, m)_{m\geq 3}, \Box, \boxslash\big\}$ and $\star' \in \{B, \overleftrightarrow{B}\}$, or $B_{\boxslash\wedge\Box}$ diagrams. In detail, if Γ is a $B_{\Box,B}$ diagram, apply (1) (c) (ii). If Γ is a $B_{\boxslash,B}$ diagram, mutate in a_1 to get a $B_{\Box,B}$ diagram of width zero. Finally, if Γ is a $B_{\perp,B}$ diagram, then mutation in a_1 allows to choose between the sub-cases of Fact 4.5 (1) (c) (iv), hence we may assume to be in sub-case (1) (c) (iv) (D). Now that we have restricted to $B_{\star\wedge\star'}$ diagrams, where $\star \in \big\{(\bigcirc, m)_{m\geq 3}, \Box, \boxslash\big\}$ and $\star' \in \{B, \overleftrightarrow{B}\}$, or $B_{\boxslash\wedge\Box}$ diagrams, Fact 4.5 (3) (b) reduces to the case of Γ being a $B_{(\bigcirc,m)\wedge\star}$ diagram, where $\star \in \{B, \overleftrightarrow{B}\}$. Once more in detail, use (3) (b) (i) (A) if Γ is a $B_{\Box\wedge B}$ diagram, (3) (b) (i) (D) if Γ is a $B_{\boxslash\wedge\Box}$ diagram and (3) (b) (iii) (A) to mutate any $B_{\boxslash\wedge B}$ diagram into a $B_{\Box\wedge B}$ diagram. Finally we are dealing with $B_{(\bigcirc,m)\wedge\star}$ diagrams with $\star \in \{B, \overleftrightarrow{B}\}$ only. Now use Fact 4.5 (2) (b) to exclude $\star = \overleftrightarrow{B}$.

Thus we may assume that Γ is a $B_{(\bigcirc,m)\wedge B}$ diagram, $m \leq |\Gamma_0| - 2$. Set $n := |\Gamma_0|$ and apply Fact 4.5 (2) (c) until Γ is a $B_{(\bigcirc,n-1)\wedge B}$ diagram. Using (2) (f) (ii) (A) of the same Fact, we are in the situation of Lemma 4.4 with Γ' a diagram with underlying undirected graph B_2, having y as an end vertex. Thus Γ is of mutation type $\mathbb{B}^{(1)}$. ☐

4.5. Type $\mathbb{C}^{(1)}$.

Fact 4.6. The following is an easy exercise in diagram mutation:

1. (1) Let Γ be a $C_{B,B}$ diagram.
   - (a) If $w(\Gamma) > 0$, then $\mu_k(\Gamma)$ is a $C_{B,B}$ diagram and $|w(\mu_k(\Gamma)) - w(\Gamma)| \leq 1$.
   - (b) Else $w(\Gamma) = 0$ and x, x' have a common neighbour $\bullet$. For k any vertex, $\mu_k(\Gamma)$ is a
     - (i) $C_{B\wedge B}$ diagram if $k = \bullet$ and $\{x, \bullet, x'\}$ carries linear orientation.
     - (ii) $C_{B,B}$ diagram else.
2. (2) Let Γ be a $C_{B\wedge B}$ diagram, k any vertex of Γ. Then $\mu_k(\Gamma)$ is a
   - (a) $C_{B\wedge B}$ diagram if $k \neq \bullet$.
   - (b) $C_{B,B}$ diagram of width zero else.

Theorem. 1.1 (5) *Let Γ be a connected diagram. Then Γ is of mutation type $\mathbb{C}^{(1)}$ if and only if it is a $C_{B,B}$ or $C_{B\wedge B}$ diagram.*

Proof. Since any diagram with underlying undirected graph of Dynkin type $\mathbb{C}$ is itself an $C_{B,B}$ diagram, one direction follows from Fact 4.6. For the other direction, we proceed by induction on $|\Gamma_0|$.

For $|\Gamma_0| = 3$ note that Fact 4.6 (2) (b) reduces to the case where Γ is a $C_{B,B}$ diagram, which has underlying graph $C_2^{(1)}$ then. Assume that Γ has $n+1$ vertices and that the statement is true for n. If Γ does not have underlying graph $C_n^{(1)}$ already, let $y \in \Gamma_0 \backslash \{x, x'\}$ such that the induced sub-graph on $\Gamma_0 \backslash \{y\}$ is connected. The rest is analogous to the proof for types $\mathbb{A}$ and $\mathbb{B}$. ☐

4.6. Type $\mathbb{D}^{(1)}$.

As for Fact 4.5, the only difficulty for the following statement lies in keeping track of notations.

Fact 4.7. The following is a straightforward exercise in diagram mutation (cf. Definition 3.6 for notations).

1. (1) Let Γ be a $D_{\star,\star'}$ diagram, where $\star, \star' \in \big\{(\bigcirc, n)_{n\geq 3}, \square, \boxslash, \perp\big\}$. Recall that the endpoints of ω_Γ are labeled x_i for some i and x'_j for some j. Let Γ_1 denote the connected component of the sub-graph of Γ induced on $\Gamma_0 \setminus \star'_0$ which is uniquely determined by containing x_i. Then Γ_1 is a $D_\star$ diagram and a full connected sub-diagram of Γ. For k any vertex of Γ we assume without loss of generality that $k \in (\Gamma_1)_0$. Then $\mu_k(\Gamma_1)$ is a $D_{\star''}$ diagram for some $\star'' \in \big\{(\bigcirc, n)_{n\geq n}, \square, \boxslash, \perp\big\}$ by Fact 4.3 and the following is true:
   - (a) if $w(\Gamma) > 0$, then $\mu_k(\Gamma)$ is a $D_{\star'',\star'}$ diagram and $|w(\mu_k(\Gamma)) - w(\Gamma)| \leq 1$.
   - (b) if $w(\Gamma) = 0$ and $k \neq x_i$, then $\mu_k(\Gamma)$ is a $D_{\star'',\star'}$ diagram and $w(\Gamma) \leq w(\mu_k(\Gamma)) \leq w(\Gamma) + 1$.
   - (c) otherwise $w(\Gamma) = 0$ and $k = x_i$. Assume that $\star$ equals
     - (i) $(\bigcirc, n)$. Then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,n+1)\vee(\bigcirc,m+1)}$ diagram if $\star' = (\bigcirc, m)$ and a $D_{(\bigcirc,n+1)\vee\star'}$ diagram else.
     - (ii) $\square$. Then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,3),(\bigcirc,n)}$ diagram of width zero if $\star' = (\bigcirc, n)$ and a $D_{(\bigcirc,3)\vee\star'}$ diagram else.
     - (iii) $\boxslash$. Then $\mu_k(\Gamma)$ is a

        (A) $D_{\boxslash\vee\star'}$ diagram if $\{x', x_1, x_2\}$ carries non-linear orientation.
        (B) $D_{\boxslash,\star'}$ diagram of width zero if $\{x', x_1, x_2\}$ carries linear orientation and $\deg(x) = 5$.
        (C) $D_{\bot,\star'}$ diagram of width zero else
    (iv) $\bot$. Then $\mu_k(\Gamma)$ is a
        (A) $D_{\boxslash\vee\star'}$ diagram if both $\{x', x, a_i\}_{1\leq i\leq 2}$ carry linear orientation.
        (B) $D_{\boxslash,\star'}$ diagram of width zero if both $\{x', x, a_i\}_{1\leq i\leq 2}$ carry non-linear orientation and $\deg(x) = 5$.
        (C) $D_{\bot,\star'}$ diagram of width zero if both $\{x', x, a_i\}_{1\leq i\leq 2}$ carry non-linear orientation and $\deg(x) = 4$.
        (D) $D_{(\bigcirc,3)\vee\star'}$ diagram else.

(2) Let Γ be a $D_{\star\vee\star'}$ diagram, where $\star, \star' \in \big\{(\bigcirc, n)_{n\geq n}, \square, \boxslash, \bot\big\}$, k a vertex of Γ.
  (a) If $k \neq \bullet$ then for
    (i) $(\star \vee \star') \neq ((\bigcirc, n) \vee (\bigcirc, m))$ we may assume without loss of generality that $k \notin \star'_0$. Denote by Γ_1 the connected component of the sub-graph of Γ induced on $\Gamma_0 \setminus \star'_0$ containing k, so that Γ_1 is a $D_\star$ diagram. Then $\mu_k(\Gamma_1)$ is a $D_{\star''}$ diagram for some $\star'' \in \big\{(\bigcirc, n)_{n\geq n}, \square, \boxslash, \bot\big\}$ by Fact 4.3. Moreover, $\mu_k(\Gamma)$ is either one of the following
        (A) a $D_{\star'',\star'}$ diagram of width zero.
        (B) or a $D_{\star''\vee\star'}$ diagram.
    (ii) $(\star \vee \star') = ((\bigcirc, n) \vee (\bigcirc, m))$ and $k \in \{a_1, a_3\}$, $\mu_k(\Gamma)$ is a $D_{(\bigcirc,m+1)\wedge\square}$ diagram if $n = 3$ and a $D_{(\bigcirc,n-1)\vee(\bigcirc,m+1)}$ diagram else.
  (b) Else $k = \bullet$. Assume that $(\star \vee \star')$ equals
    (i) $((\bigcirc, 3)\vee(\bigcirc, 3))$. Then $\mu_k(\Gamma)$ is a $D_{\square\vee\square}$ diagram if both $x_3, x'_3 \in \Gamma_0$ and a $D_{\square\vee\bot}$ diagram else.
    (ii) $((\bigcirc, n) \vee (\bigcirc, 3))$ with $n \neq 3$. Then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,n-1)\vee\square}$ if $x'_3 \in \Gamma_0$ and a $D_{(\bigcirc,n-1)\vee\bot}$ diagram else.
    (iii) $((\bigcirc, n) \vee (\bigcirc, m))$ with $(n, m) \neq (3, 3)$. Then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,n-1),(\bigcirc,m-1)}$ diagram of width zero.
    (iv) $(\bigcirc, n) \vee \boxslash$. Then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,n+1)\wedge\boxslash}$ diagram.
    (v) $(\bigcirc, n) \vee \square$. Then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,n+1)\wedge(\bigcirc,3)}$ diagram.
    (vi) $(\bigcirc, n) \vee \bot$. Then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,n+1)\wedge\boxslash}$ diagram if $\{a'_1, a'_2, \bullet\}$ carries non-linear orientation and a $D_{(\bigcirc,n+1)\wedge(\bigcirc,3)}$ diagram else.
    (vii) $(\boxslash \vee \boxslash)$. Then assuming without loss of generality that $x_2 = \bullet = x'_1$, $\mu_k(\Gamma)$ is a $D_{\boxslash\vee\boxslash}$ diagram if $\{x_1, \bullet, x'_2\}$ carries non-linear orientation and a $D_{\boxtimes}$ diagram else.
    (viii) $(\boxslash \vee \square)$. Then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,3)\wedge\boxslash}$ diagram.
    (ix) $(\square \vee \square)$. Then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,3)\vee(\bigcirc,3)}$ diagram.
    (x) $(\boxslash \vee \bot)$. Then $\mu_k(\Gamma)$ is a $D_{\boxslash\vee\bot}$ diagram if one of $\deg^\pm(\bullet)$ equals four. Else one of $\deg^\pm(\bullet)$ equals three and μ_k yields a $D_{\boxtimes}$ diagram if $\{a'_1, a'_2, \bullet\}$ carries non-linear orientation and $D_{(\bigcirc,3)\wedge\boxslash}$ diagram else.

    (xi) $(\square \vee \perp)$. Then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,3)\wedge\boxslash}$ diagram if $\{a'_1, a'_2, \bullet\}$ carries non-linear orientation and $D_{(\bigcirc,3)\vee(\bigcirc,3)}$ diagram else.
    (xii) $(\perp \vee \perp)$. Then $\mu_k(\Gamma)$ is a $D_{\perp\vee\perp}$ diagram if one of $\deg^{\pm}(\bullet)$ equals four, a $D_{(\bigcirc,3)\wedge\boxslash}$ diagram if one of $\deg^{\pm}(\bullet)$ equals three and a $D_{\boxtimes}$ diagram else.

(3) Let Γ be a $D_{(\bigcirc,n)\wedge\star}$ diagram, where $\star \in \{\square, \boxslash, \overleftrightarrow{\boxslash}\}$.
  (a) If $k \notin ((\bigcirc, n) \wedge \star)_0$ then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,n)\wedge\star}$ diagram.
  (b) If $k = x_i$ for some i then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,n+1)\wedge\star'}$ diagram.
  (c) If $k \in ((\bigcirc, n) \wedge \star)_0 \backslash \{\bullet_j, a'_j\}_{1\leq j\leq 2}$ and $n > 3$, then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,n-1)\wedge\star}$ diagram.
  (d) If $k \in ((\bigcirc, 3) \wedge \star)_0 \setminus \{\bullet_j, a'_j\}_{1\leq j\leq 2}$, then $\mu_k(\Gamma)$ is a $D_{\boxslash\wedge\boxslash}$ diagram if $\star = \overleftrightarrow{\boxslash}$. Else $\star = \boxslash$ and μ_k yields a $D_{\square\wedge\square}$ diagram.
  (e) Else $k \in \{\bullet_j, a'_j\}_{1\leq j\leq 2}$. Assume $\star$ equals
    (i) $\overleftrightarrow{\boxslash}$. If $k \in \{\bullet_1, \bullet_2\}$ then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,n)\wedge\overleftrightarrow{\boxslash}}$ diagram. Else $k \in \{a'_1, a'_2\}$ and μ_k yields a $D_{(\bigcirc,n+1)\wedge\square}$ diagram.
    (ii) $\boxslash$. If $k \in \{a'_1, a'_2\}$ then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,n+1)\wedge\square}$ diagram. Else $k \in \{\bullet_1, \bullet_2\}$ and
      (A) for $n > 3$, $\mu_k(\Gamma)$ is a $D_{(\bigcirc,n-1)\vee\boxslash}$ diagram of width zero if $\deg(k) = 5$ and a $D_{(\bigcirc,n-1)\vee\perp}$ of width zero else.
      (B) for $n = 3$, $\mu_k(\Gamma)$ is a $D_{\boxslash\wedge\square}$ diagram if $\{x_1, x_2, a'_1\} \subset \Gamma$. Else $|\ \{x_1, x_2, a'_1\} \cap \Gamma\ | = 2$ and $\mu_k(\Gamma)$ is a
        (α) $D_{\boxslash\vee\perp}$ of width zero if $\deg(k) = 5$.
        (β) $D_{\square\vee\perp}$ of width zero else.
    (iii) $\square$.
      (A) If $n > 4$ then $\mu_k(\Gamma)$ is a
        (α) $D_{(\bigcirc,n-1)\vee(\bigcirc,3)}$ diagram if $k \in \{\bullet_1, \bullet_2\}$. Moreover if $\deg(k) = 3$ then $x'_3 \notin (\mu_k(\Gamma))_0$.
        (β) Else $k \in \{a'_1, a'_2\}$ and $\mu_{a'_1}(\Gamma)$ is a $D_{(\bigcirc,n-1)\wedge\boxslash}$ diagram while $\mu_{a'_2}(\Gamma)$ is a $D_{(\bigcirc,n-1)\wedge\overleftrightarrow{\boxslash}}$ diagram.
      (B) If $n = 3$ then $\mu_k(\Gamma)$ is a
        (α) $D_{(\bigcirc,3)\wedge\square}$ diagram if $k \in \{\bullet_1, \bullet_2\}$.
        (β) Else $k \in \{a'_1, a'_2\}$ and $\mu_{a'_1}(\Gamma)$ is a $D_{\square\wedge\square}$ diagram while $\mu_{a'_2}(\Gamma)$ is a $D_{\boxslash\wedge\boxslash}$ diagram.

(4) Let Γ be a $D_{\star\wedge\star}$ diagram, where $\star \in \{\square, \boxslash\}$.
  (a) If $k \notin (\star \wedge \star)_0$ then $\mu_k(\Gamma)$ is a $D_{\star\wedge\star}$ diagram.
  (b) Else $k \in (\star \wedge \star)_0$.
    (i) If $k = \bullet$ then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,3)\wedge\boxslash}$ diagram if $\star = \square$ and $D_{(\bigcirc,3)\wedge\overleftrightarrow{\boxslash}}$ diagram else.
    (ii) If $k \in \{x_1, x_2\}$ then $\mu_k(\Gamma)$ is a $D_{\boxtimes}$ diagram if $\star = \square$ and a $D_{\boxslash\wedge\boxslash}$ diagram else.
    (iii) If $k \in \{a_1, a_2\}$ then $\mu_k(\Gamma)$ is a $D_{(\bigcirc,3)\wedge\square}$ diagram.

(5) Let Γ be a $D_{\boxtimes}$ diagram.

(a) If $k \notin \boxtimes_0$ then $\mu_k(\Gamma)$ is a $D_{\boxtimes}$ diagram.

Else $k \in \boxtimes_0$.
(b) If $k \in \{a_i\}_{1 \leq i \leq 4}$ then $\mu_k(\Gamma)$ is a $D_{\square \wedge \square}$ diagram.
(c) If $k = x_1$ then $\mu_k(\Gamma)$ is a
  (i) $D_{\boxslash \vee \boxslash}$ diagram if $\deg(x) = 6$.
  (ii) $D_{\boxslash \vee \perp}$ diagram if $\deg(x) = 5$.
  (iii) $D_{\perp \vee \perp}$ diagram else.

The last part of Theorem 1.1 is proved in the same way as part (4). The difficult part is to show that any diagram belonging to one of the families defined in subsection 3.6 is of mutation type $\mathbb{D}$. This is done by excluding one by one the cases to be considered, the essential point being that we have a complete overview of the effect of mutation on these diagrams by Fact 4.7.

Theorem. 1.1 (6) *Let* Γ *be a connected diagram. Then* Γ *is of mutation type* $\mathbb{D}^{(1)}$ *if and only if it is a* $D_{\star,\star'}, D_{\star\vee\star'}, D_{\star\wedge\star'}$ *diagram, where* $\star, \star' \in \left\{(\bigcirc, n)_{n\geq 3}, \square, \boxslash, \perp, \overleftrightarrow{\boxslash}\right\}$, *or a* $D_{\boxtimes}$ *diagram.*

Proof. Any diagram with underlying graph of Dynkin type $\mathbb{D}^{(1)}$ is a $D_{\perp,\perp}$ diagram, hence Fact 4.7 implies that any diagram of mutation type $\mathbb{D}^{(1)}$ is a $D_{\star,\star'}, D_{\star\vee\star'}, D_{\star\wedge\star'}$ diagram, where $\star, \star' \in \left\{(\bigcirc, n)_{n\geq 3}, \square, \boxslash, \perp, \overleftrightarrow{\boxslash}\right\}$, or a $D_{\boxtimes}$ diagram.

To show the other direction, first assume that $|\Gamma_0| = 5$. This restricts the possibilities for Γ to $D_{\star\wedge\star'}$, where $\star, \star' \in \left\{(\bigcirc, 3), \square, \boxslash, \overleftrightarrow{\boxslash}\right\}$, or $D_{\boxtimes}$, or $D_{\perp\vee\perp}$ diagrams. Note that the last mentioned has underlying undirected graph $D_4^{(1)}$ already. Moreover Fact 4.7 (5) (c) (iii) implies that $D_{\boxtimes}$ diagrams are also of mutation type $\mathbb{D}^{(1)}$. Hence we only have to deal with $D_{\star\wedge\star'}$ diagrams, where $\star, \star' \in \left\{(\bigcirc, 3), \square, \boxslash, \overleftrightarrow{\boxslash}\right\}$. By Fact 4.7 (4) (b) (iii) we may restrict to $D_{(\bigcirc,3)\wedge\star}$ diagrams, where $\star \in \{\square, \boxslash, \overleftrightarrow{\boxslash}\}$. Now (3) (e) (i) and (ii) of the same Fact imply that it suffices to deal with $D_{(\bigcirc,3)\wedge\square}$ diagrams. Using (3) (e) (iii) (B) (β) we see that we may restrict to $D_{\square\wedge\square}$ diagrams after all. But then (4) (b) (ii) yields that all these diagrams are mutation equivalent to $D_{\boxtimes}$ diagrams, which we already know to be of mutation type $\mathbb{D}^{(1)}$. This finishes the proof for $|\Gamma_0| = 5$

For $|\Gamma_0| > 5$ we apply the results of Fact 4.7 in order to reduce to the case where Γ is a $D_{(\bigcirc,m)\wedge\square}$ diagram, where $m \leq |\Gamma_0| - 2$. First note that any $D_{\star,\star'}$ diagram, where $\star, \star' \in \left\{(\bigcirc, m)_{m\geq 3}, \square, \boxslash, \perp\right\}$, is mutation equivalent to a $D_{\bar{\star},\bar{\star}'}$ diagram, where $\bar{\star}, \bar{\star}' \in \left\{(\bigcirc, m)_{m\geq 3}, \square, \boxslash, \perp\right\}$, of width zero. Hence (1) (c) and (5) (b) reduce to the case of $D_{\star\vee\star'}, D_{\star\wedge\star'}$ diagrams with $\star, \star' \in \left\{(\bigcirc, m)_{m\geq 3}, \square, \boxslash, \perp, \overleftrightarrow{\boxslash}\right\}$; here we use that for $D_{\boxslash,\star'}$ and $D_{\perp,\star}$ diagrams of width zero we may, by subsequent mutation in a_1 and a_2, always assume to be in case (1) (c) (iii) (A) respectively (1) (c) (iv) (A) of Fact 4.7. Next, (2) (b) reduces to $D_{\star\wedge\star'}$ diagrams with $\star, \star' \in \left\{(\bigcirc, m)_{m\geq 3}, \square, \boxslash, \overleftrightarrow{\boxslash}\right\}$: this is immediate for $(\bigcirc, n) \vee \star'$, $\star' \neq (\bigcirc, m)$, and $(\boxslash \vee \square)$. For $(\star \vee \perp)$, $\star \in \{\square, \boxslash, \perp\}$, use that subsequent mutation in a_1 and a_2 allows for choosing any of the cases (2) (b) (x), (xi) or (xii). If $(\star \vee \star') \in \{(\square \vee \square), (\boxslash \vee \boxslash)\}$ then mutation in a_1 yields a $D_{(\boxslash\vee\square)}$ diagram. For $(\star \vee \star') = ((\bigcirc, n), (\bigcirc, m))$ use (2) (a) (ii) $n - 2$ times. Hence we have reduced to $D_{\star\wedge\star'}$ diagrams, where $\star, \star' \in \left\{(\bigcirc, m)_{m\geq 3}, \square, \boxslash, \overleftrightarrow{\boxslash}\right\}$. Now apply Fact 4.7 (4) (b) (iii) to exclude $\star, \star' \in \{\boxslash, \square\}$ and (3) (e) (i) as well as (ii) to get rid of $\star \wedge \star' \in \left\{(\bigcirc, m) \wedge \boxslash, (\bigcirc, m) \wedge \overleftrightarrow{\boxslash}\right\}$.

Thus we may assume that Γ is a $D_{(\bigcirc,m)\wedge\square}$ diagram, where $m \leq |\Gamma_0| - 2$. Set $n := |\Gamma_0|$ and apply Fact 4.7 (3) (b) until Γ is a $D_{(\bigcirc,n-2)\wedge\square}$ diagram. Due to cardinality Fact 4.7 (3) (e) (iii) (A) (α) yields that we may assume Γ to be of the form as described in the second case of (2) (b) (ii) of the same Fact; hence we are in the situation of Lemma 4.4 with Γ' a diagram with underlying undirected graph A_3, having y as its middle vertex. Thus Γ is of mutation type $\mathbb{D}^{(1)}$. $\square$

References

1. M. Barot, C. Geiss, and A. Zelevinsky, *Cluster algebras of finite type and positive symmetrizable matrices*, J. Lon. Math. Soc. **73 (3)** (2006), 545–564.
2. J. Bastian, *Mutation classes of $\tilde{A}_n-$ quivers and derived equivalence classification of cluster tilted algebras of type $\tilde{A}_n$*, arXiv:0901.1515.
3. A. B. Buan, R. Marsh, M. Reineke, I. Reiten, and G. Todorov, *Tilting theory and cluster combinatorics*, Adv. Math. **204 (2)** (2006), 572–618.
4. A. B. Buan, R. Marsh, and I. Reiten, *Cluster-tilted algebras*, Trans. Amer. Math. Soc. **359 (1)** (2007), 323–332.
5. A. B. Buan and I. Reiten, *Acyclic quivers of finite mutation type*, International Math. Research Notices2006, Art. ID12804.
6. A. B. Buan, I. Reiten, and A. Seven, *Tame concealed algebras and cluster quivers of minimal infinite type*, J. of Pure and Applied Algebra **211** (2007), 71–82.
7. A. B. Buan and D. F. Vatne, *Derived equivalence classification for cluster-tilted algebras of type A_n*, J. Algebra **319 (7)** (2008), 2723–2738.
8. S. Fomin and A. Zelevinsky, *Cluster algebras i. foundations*, J. Amer. Math. Soc. **15 (2)** (2002), 497–529.
9. ______, *Cluster algebras ii. finite type classification*, Invent. Math **154 (1)** (2003), 63–121.
10. W. Ge, H. Lv, and S. Zhang, *Cluster-tilted algebras of type D_n*, arXiv:0812.0650.
11. D. Happel and D. Vossieck, *Minimal algebras of infinite representation type with preprojective component*, Manuscripta Math. **42 (2-3)** (1983), 221–243.
12. B. Keller, http://www.math.jussieu.fr/∼keller/quivermutation/.
13. ______, *Cluster algebras, quiver representations and triangulated categories*, arXiv:0807.1960.
14. R. Schiffler, *A geometric model for cluster category of type D_n*, J. Algebra Com. **27 (1)** (2008), 1–21.
15. A. Seven, *Recognizing cluster algebras of finite type*, The Electronic Journal of Combinatorics **14** (2007).
16. D. Vatne, *The mutation class of D_n quivers*, arXiv:0810.4789.

MATHEMATISCHES INSTITUT DER UNIVERSITÄT BONN, ENDENICHER ALLEE 60, D-53115 BONN.

E-mail address: `henrich@math.uni-bonn.de`